



“Sexual health is a state of *physical, emotional, mental and social well-being* in relation to sexuality; it is not merely the absence of disease, dysfunction or infirmity. Sexual health requires a positive and respectful approach to sexuality and sexual relationships, as well as the possibility of having pleasurable and safe sexual experiences, free of coercion, discrimination and violence”

Female Sexual Dysfunction: Screening, Diagnosis and Treatment



**MARY LYNN, DO
ASSISTANT PROFESSOR
DEPARTMENT OF OBSTETRICS AND GYNECOLOGY**

**CO-DIRECTOR OF THE SEXUAL WELLNESS CLINIC
LOYOLA UNIVERSITY MEDICAL CENTER**

Goals and Objectives



- Identify barriers to screening for female sexual dysfunction
- Identify the most prevalent types of female sexual dysfunction
- Discuss diagnostic considerations for female sexual dysfunction and when to begin treatment
- Discuss strategies for female sexual dysfunction

Prevalence of Sexual Dysfunctions




Men N > 90,000	Women N ~ 10,000
<ul style="list-style-type: none">■ Any SD 31%■ Low desire 5% - 15%■ Anorgasmic 8%■ Rapid ejaculation 14% - 30%■ Erectile disorder(ED) 18% - 52%■ 36% moderate or complete ED■ ↑ with age -<ul style="list-style-type: none">- 25% of men < 59 yo- 61% of men > 70	<ul style="list-style-type: none">■ Any SD 32%■ Low desire 17% - 55%<ul style="list-style-type: none">> in surgical menopause■ Arousal problems 14% - 35%■ Orgasm issues 25% -39%■ Pain 2% - 26%

- ❖ Sexual problems can be life long or acquired
 - ❖ Generalized or situational

Impact of Sexual Problems



- Global Study of Sexual Attitudes Behaviors



- Survey of 27,000+, 40-80 y/o

- 80% of men, 60% of women rated sexual function as a “**moderately to extremely**” important

- Not all pts meet formal diagnostic criteria but the impact of sexual problems is ↑
- Treatment helpful even if a formal dx is not made

Correlates of Sexual Problems



- **Age**
 - Occurrence of problems ↑ with age
 - Distress ↓ with age
- **Health comorbidities**
 - DM, CVD, HTN, prostate / gyn problems, cancer, obesity, tobacco, alcohol, recreational drug use
- **Medications**
 - Psychotropics, antihypertensives, anticonvulsants..
 - ~ 25% of ED cases are medication related
- **Psychosocial factors**
 - Depr, anx, relationship issues, stress, sexual trauma

Emotional Factors and Sexuality



- **Depression**
 - Impacts all aspects of sexual response
 - ✦ desire, arousal and orgasm
- **Anxiety**
 - Common in SD - performance anxiety, fear of inadequacy, spectating
 - ✦ all impede psychophysiological arousal
- **Anger**
 - Impedes communication and intimacy

Depression and Sexual Dysfunction



- Sexual dysfunction is both:
 - A symptom of depression
 - An adverse effect of many antidepressants & other psychotropics
- Treatment-emergent sexual dysfunction
 - A major cause of noncompliance and drug discontinuation
 - It is a substantial risk factor for relapse or recurrence of a depressive episode
 - Important to assess sexual function in patients with depression before selecting the most appropriate antidepressant medication

Importance of Screening for SD



Underdiagnosed and Undertreated

- Obstacles for patients
 - ✦ Don't ask, don't tell..
 - ✦ Patients not likely to bring it up unless asked
- Obstacles for physicians
 - ✦ Lack of training
 - ✦ Lack of confidence
 - ✦ Lack of knowledge regarding treatment options
 - ✦ Inadequate time to obtain a sexual history
 - ✦ Underestimation of the prevalence of sexual dysfunction

Easy Screening Questions



- Are you sexually active?
- Any pain during sexual activity?
- Are you able to achieve an orgasm?
- Any decreased desire or libido that is troubling for you and your partner?
- Everyone should be screened for domestic violence and sexual abuse during annual visits.

Brief Sexual Symptom Checklist



1. *Are you satisfied with your sexual function?* Yes No

If No, please continue.

2. *How long have you been dissatisfied with your sexual function?*

3a. *men/women specific questions...*

3b. *Which problem is most bothersome (circle)* 1 2 3 4 5 6 7

4. *Would you like to talk about it with your doctor?*

Yes No

Brief Sexual Symptom Checklist: 3a



For Men (BSSC-M)

3a. The problems with your sexual function is: (mark one or more)

- 1 Problems with little or no interest in sex**
- 2 Problems with erection**
- 3 Problems ejaculating too early during sexual activity**
- 4 Problems taking too long, or not being able to ejaculate or have orgasm**
- 5 Problems with pain during sex**
- 6 Problems with penile curvature during erection**
- 7 Other :**

For Women (BSSC-W)

3a. The problems with your sexual function is: (mark one or more)

- 1 Problems with little or no interest in sex**
- 2 Problems with decreased genital sensation (feeling)**
- 3 Problems with decreased vaginal lubrication (dryness)**
- 4 Problems reaching orgasm**
- 5 Problems with pain during sex**
- 6 Other :**

Integrative Treatment Models



- **PLISSIT Model**

(Annon, 1976)

- **Permission** – for the pt to discuss the issue
- **Limited Information** – education about the psychophysiology of sexual arousal and normal sexual functioning
- **Specific Suggestions** – e.g., communication skills, relaxation skills, sensate focus
- **Intensive Therapy** - refer for additional treatment as needed

- **Brief Sexual Counseling**

(Schover & Jensen, 1988)

- Sex education
- Restructure maladaptive beliefs about sexuality
- Help the pt stay sexually active (e.g., sensate focus, identify mutually satisfying sexual experiences)
- Address conflict resolution and communication skills

Sex Therapy – Patient Centered



- Starts with psychosexual evaluation:
 - Assessment of the pt's / couple's PP and sexual history
 - Current sexual practices
 - Relationship quality
 - Emotional health
 - Contextual factors (e.g., chr illness, stressors, etc)
 - Psychosexual & developmental history
 - Review of relevant medical and biological factors
 - Physical exam / sexological exam
 - Mutual goal setting by pt/partners

Treatment Components



- ❑ Body image exercises
- ❑ Sensate focus
 - ❑ 5-step exercise for sensual/sexual intimate touch
- ❑ Anxiety management
 - ❑ Relaxation training, systematic desensitization to feared stimuli
- ❑ Bibliotherapy - erotic reading
 - ❑ The Busy Couples Guide to Great Sex (McAllister & Rallie)
 - ❑ The Art of Kissing (Cane)
 - ❑ Fantasy reading

Female Sexual Interest/Arousal Disorder



- At least 3 of the following:
 - Reduced (or absent) interest in sexual activity
 - Reduced sexual/erotic thoughts or fantasies
 - Reduced initiation of sexual activity, unreceptive to a partner's attempts
 - Reduced sexual excitement or pleasure
 - Reduced sexual interest/arousal in response to sexual cues (e.g., written, verbal, visual)
 - Reduced genital or nongenital sensations
- Freq associated with:
 - Orgasm problems, painful sex, couple-level discrepancies in desire, unrealistic expectations, lack of information about sexuality

© Original Artist / Search ID: abrn192




Rights Available from CartoonStock.com

"I really think you should see a specialist about your lack of libido Sharon."

Basson's Non-linear Model



FIGURE 3. Non-linear Model of Female Sexual Response Developed by Basson⁶



Basson's non-linear model acknowledges how emotional intimacy, sexual stimuli, and relationship satisfaction affect female sexual response.

Treatment



- Biopsychosocial and Educational Approach
 - Focus on the Cause of the Disorder
 - Cognitive-Behavioral Techniques and/or Traditional Sex Therapy.
 - ✦ Communication exercises
 - ✦ Body image exercises
 - ✦ Sensate focus exercises
 - Mindfulness-based treatment
 - ✦ Encouraging results
 - ✦ Need larger studies
- Pharmacological treatment

Check Medications



- Medications commonly associated with SD
 - Antihypertensives
 - Histamine blockers
 - Oral Contraceptive pills
 - Psychotropic medications
 - ✦ SSRIs are most commonly linked to sexual dysfunction
 - ✦ Estimated incidence of SSRI-induced sexual dysfunction ranges from approximately 15 to 80 percent
 - ✦ Interest/Arousal and Orgasmic disorder are the most common issues

Meds cont..



- Decreasing the dosage may help alleviate some issues
- Switching to another antidepressant may alleviate symptoms, but other antidepressant classes also have been associated with sexual dysfunction.
- A structured treatment interruption may be helpful in some patients, but is not always an option in some other patients.

Flibanserin



- 5-HT serotonin receptor agonist and a dopamine D4 receptor partial agonist.
- Non-Hormonal
- Increases dopamine/noradrenalin and reduces Serotonin
- Hypoactive sexual desire disorder (HSDD) in premenopausal women
- Dose 100mg qhs
- Side effects: fatigue (morning), hypotension
- Take daily and no alcohol use on this medication.

Female Orgasmic Disorder



Diagnostic Criteria – for at least 6 months


- Either of the following on 75%–100% during sexual activity:
 - Marked delay in, marked infrequency of, or absence of orgasm.
 - Markedly reduced intensity of orgasmic sensations.
- Causes clinically significant distress
- Not explained by
 - A nonsexual mental disorder
 - A consequence of severe relationship distress (e.g., partner violence)
 - Other significant stressors
 - Not due to the effects of a substance/medication
 - Not due to another medical condition.

Female Orgasmic Disorder




- Timeline to relating to Orgasmic disorder
 - Ever had an orgasm?
 - Had one with this relationship?
 - Does partner know?
 - Does she engage in self exploration?
- Any comorbid factors (new or old)

Males



Females





- **75% of all males → orgasm is possible within the first 4 minutes after initiation of sexual intercourse**
- **All women the average time to reach orgasm is between 10 and 20 minutes**

Treatment:



- Focus on being comfortable your body and self exploration is encouraged.
- Directed self exploration exercises with clitoral stimulation:
 - encourage patience and persistence with at least three weekly sessions in a good setting
- Transfer of self exercises to “couple”
 - Allow partner to first observe then engage in self exploration exercises. Consider your sexual needs first over your partners.
- Bibliotherapy (see erotic reading list also)
 - The G spot or The science of orgasm
- Lubricants
 - Zestra – stimulating gel (otc)

Genito-Pelvic Pain/Penetration Disorder



- Sxs highly comorbid (need 1 of 4 to dx)
 - Difficulty having intercourse
 - Genito-pelvic pain
 - Fear of pain or vaginal penetration
 - Tension of the pelvic floor muscles
- Behavioral avoidance of sexual situations and of gyn exams is common
 - Avoidance pattern is similar to phobic disorders

Causes of Dyspareunia



- **Atrophy**

- Leading cause of dyspareunia due to decreased estrogen

- Causes:

- ✦ Menopause
- ✦ Premature Ovarian Failure
- ✦ Hypothalamic Amenorrhea (excessive exercise or rapid weight loss)
- ✦ Postpartum/Breastfeeding
- ✦ Low Estrogen Contraceptives
- ✦ Radiation or Chemotherapy (Tamoxifen).

Atrophy



Treatment:

- ✦ Hormone-free Lubricants (water-base or silicon):
 - With intercourse
 - Free of Parabens and Glycerin
- ✦ Hormone-free Moisturizers
 - Every third night
- ✦ Local estrogen therapy: cream, tablet or vaginal ring.
 - If history of breast cancer – discuss with oncologist prior to use.
- ✦ Pelvic floor physical therapy (dilators if necessary)
- ✦ Relaxation training.

Vaginismus



- Prevalence rates ranging from 1% to 6%
- Cannot consummate intercourse because vaginal penetration is not possible
 - Involuntary spasm of perineal/levator muscles
 - Vaginal muscle contractions occur as an automatic defense to vaginal penetration
 - For some women it is only limited to vaginal exams, but intercourse is possible and comfortable.
- Diagnosed by eliciting muscle spasm by depressing the levators

Treatments cont..



- Relaxation and desensitization techniques
 - Deep muscle relaxation techniques to use during exercises
 - Using dilators
 - ✦ Starting with the smallest one that is comfortable
 - ✦ Gradually over time increasing diameter of the dilator as tolerated.
 - ✦ Goal is to desensitize a woman to her fear that vaginal penetration will be painful
 - ✦ Enable her to gain a sense of control over a sexual encounter or a pelvic examination
- Pelvic floor physical therapy
- Vaginal valium (compounded into a suppository) may be helpful.

Take Home Points



- Roughly 1/3 of patients have Sexual Dysfunction
- It is important to Screen for it Annually
- The Most Common Types of dysfunction are:
 - Hypoactive sexual desire disorder (decreased libido)
 - Anorgasmia
 - Genito-pelvic pain disorder (dyspareunia)
- Treatment will usually start with Behavioral Modifications and Education
- Pelvic Floor Physical Therapy can be very helpful for Dyspareunia.

References




- Clayton A, Pradko J, Croft H, et al. Prevalence of sexual dysfunction among newer antidepressants. *J Clin Psychiatry*. 2002;63:357-366.
- Kennedy, Sidney H. MD, FRCPC*†; Rizvi, Sakina HBSoc. Sexual Dysfunction, Depression, and the Impact of Antidepressants. *Journal of Clinical Psychopharmacology*. Lippincott Williams & Wilkins, Inc. Volume 29(2), April 2009, pp 157-164
- <http://fsfi-questionnaire.com/FSFI%20questionnaire2000.pdf>
- Hatzichristou D, Rosen RC, et al. *J Sex Med*. 2004;1:49-57
- Sensate focus Protocol adapted from Loyola Sexual Wellness Patient Handbook 2013.
- Erotic Reading list adapted from: The Pelvic & Sexual Health Institute, Philadelphia (Copyright 2006)
- Brotto LA, Petkau JA, Labrie F, Basson R: Predictors of sexual desire disorder in women. *J Sex Med* 8(3):742-753, 2011
- Cartoonstock.com

References



- Annon Tx. The behavioral treatment of sexual problems, vol 1, Bret Therapy, Enabling systems, Honolulu, 1974.
- Basson R et al. Summary of the recommendations on sexual dysfunctions in women. *J of Sexual Medicine* 2004:1:24-34.
- DeRogatis LR, & Burnett AL. The epidemiology of sexual dysfunctions. *J of Sexual Medicine* 2008:5:289-300.
- Dunn KM, Croft PR, Hackett GI. Association of sexual problems with social, psychological, and physical problems in men and women: A cross sectional population survey. *J of Epidemiol Community Health* 1999:53:144-148.
- Hatzichristou D et al. Clinical evaluation and management strategy for sexual dysfunction in men and women. *J of Sexual Medicine* 2004:1:49-57.
- Leiblum SR. Principles and practice of sex therapy (4th Ed.). NY: Guilford Press, 2007.
- Renshaw D. Seven weeks to better sex (3rd Ed.). Redondo Beach, CA: Westcom Press, 2004.

Thank You!



search ID: ccan133

"How was it for you?"